



Article

Palaeoecological Aspects of Nannoplankton Assemblages of AS-2 Well, Niger Delta

*Ekaete Enamekere Umoh

Department of Earth Sciences, Faculty of Natural Sciences, Ajayi Crowther University, Oyo, Oyo State, Nigeria.

ee.umoh@acu.edu.ng (E.E.U.)

* Correspondence: ee.umoh@acu.edu.ng; +234-8056183513

Article history: received, May 1, 2023; revised, June 7, 2023; accepted, June 9, 2023; published: June 14, 2023.

Abstract

Certain nannoplankton species are sensitive to surface water conditions hence, they are proxy for palaeoecological studies. This study analyzed biostratigraphy and inferred paleoecologic preferences of major nannofossil species of ditch cutting samples from well AS-2, Niger delta. The relative fluctuation and changes in abundance of the nannoplankton allowed the following deductions in the study interval: Two major biozones, NP21 and NP20 of late Eocene (Priabonian) and early Oligocene (Rupelian). Higher abundances of *Reticulofenestra* spp (49%) compared to the total taxa recovered, including *Cyclicargolithus* spp, *Coccolithus* spp and *Chiasmolithus* spp were characterized as cool water, eutrophic conditions while *Sphenolithus* spp, *Helicosphaera* spp, *Pontosphaera* spp were characterize as warmer water more Oligotrophic species. The total increase in abundance of cooler-water indicators over the warm-water forms within the study intervals suggest a prevailing environmental condition describing cooler water trends, eutrophic and upwelling ones with nutrient rich conditions during the Oligocene period than in the Eocene.

Keywords: Nannoplankton, Biostratigraphy, Paleoecology, Oligotrophic, Niger Delta.

1. Introduction

The term “Nannoplankton” was defined by [1] as small plankton, that will pass through the mesh of the phytoplankton net of 63microns. Calcareous nannofossils have been identified as one of the most abundant phytoplankton group in the marine realm with enormous effect to changes that occur in the oceanographic terrain. This phytoplankton group has progressively become tremendously useful in solving Biostratigraphic [2] palaeologic, palaeoceanographic, palaeoclimatic and even palaeoenvironmental problems [3 -8].

Nannoplanktons are known for their distinct, pronounced and quick response to changes to temperature, light, availability of nutrients, salinity, primary productivity and other parameters in the world oceans, all these have extensively been studied in many basins *around the world*. Thus, more efforts in research is necessary towards understanding the Mesozoic and Cenozoic marine systems and interactions as evidenced by nannofossils [3],[8-10] for classical aspects of palaeologic and palaeoceanographic interpretations even in the global and local basins.

Some vital reasons for calcareous nannofossils importance as explained by many authors include: they are one of the most abundant phytoplankton species of the marine realm; the distribution patterns of nannoplanktons species give clues to stratigraphic and oceanographic information; they are most abundant in marine sediments and show many evolutionary changes [11], and several individual taxon studied have ecologic affinity for different environments.

Review of available literature has shown that the Paleogene sections of the Niger Delta are yet to be studied in much detail as their Neogene and Pleistocene counterparts [12], [13], [14], and [15]. The studied interval which penetrates the Eocene – Oligocene is investigated for some dominant taxa of nannoplankton in sediments for their palaeocological aspects. This present study is aimed at establishing the biozonation, age and the palaeocological affinity of some selected nannoplankton recovered from the studied section in Niger Delta, and along with published catalogues identify those with significant information to bridge the information gap palaeocologically in the Nigerian local oceanographic settings. The information established will contribute conclusively that different calcareous nannofossil assemblages can be used to characterize the different palaeocological preferences in sections of the Niger Delta and the data can be applied in the recognition and interpretation of past oceanic conditions. The popular Niger Delta basin has been the major source of hydrocarbon exploration since 1957, and today it is named the Africa's leading oil province, penetrating more than 5,000 wells. The exploration activity has since shifted to the offshore part of the Niger Delta hence, more revelation by research into the oceanic setting of the different Era, period; stages will be helpful for the relatively new exploration and exploitation activities in the basin.

1.1 Location of the study area

The study area is an onshore well AS-2 (coded name) from Greater Ughelli depobelt, Niger Delta (Fig. 1). The basin is located in the Niger Delta basin which lies between longitudes 3°E and 9°E and latitudes 4°E and 5°2'N.




Figure 1: Depobelts of the Niger Delta and Location of study well (modified from [16]).

1.2 Outline Geology and Stratigraphy of Niger Delta

The Niger Delta is situated on the Gulf of Guinea at the West coast of Central Africa. The Niger-Delta is termed as the world's largest and also with sub aerial portion covered by about 75 000km² and extending more than 300km from apex to mouth, and the clastic sediments regressive wedge reaches

about 12km of maximum thickness [17]. The basin is as a result of the opening of the South Atlantic Ocean between Africa and South America continents. Development started in the late Paleocene and Eocene, these time sediments began to build out beyond troughs between basement horst blocks at the northern flank of the present day delta area and since then the delta plain has prograded southwards onto oceanic crust and also gradually assuming a convex to the sea morphology [17]. The rate of sediment supply and subsidence had controlled the structure and stratigraphy in the delta. Three formation names are globally used [18 -19] in the oil industry these three lithofacies are usually referred to as Akata, Agbada and Benin formations. Akata Formation (marine shales) is the basal unit, composed of shales, silts and clays. The approximate thickness of this sequence may be 7000m. Marine shales form the base of the sequence in each depobelts and range from paleocene and Holocene in age (Fig 2). The Agbada Formation (paralic clastics): It forms the hydrocarbon prospective sequence in the Niger delta. It is represented by an alternation of sands, clays and silts in various proportions and thickness, representing cyclic sequence of offlap units. These paralic clastics are truly deltaic sequence and were deposited in a number of delta front, delta topset and fluviodeltaic environment [17]. The alternation of coarse and fine sediment provides multiple reservoir-seal couplets. The paralic marine shale sequence is present in all depobelts, and ranges in age from the Eocene to Pleistocene (Fig 2). Benin Formation (continental sands): It is the shallowest part of the sequence and is composed entirely of non-marine sands. It was deposited in alluvial environment by a southward shift of deltaic deposition into a new depobelt. The oldest continental sands are probably Oligocene, but they lack fauna and are impossible to date directly. They become thinner offshore and disappear near the continental shelf edge (Fig 2).




Figure 2: Column of the three stratigraphic formations of Niger Delta (Modified from [17]).

2. Methods

Data used in this study include gamma ray log and ditch cuttings samples from the interval of 10510 - 13790ft from well AS-2 of Greater Ughelli depobelt, Niger Delta. Detailed analysis and interpretation of the gamma ray log in combination with petrographic studies for lithologic description and analyses for nannofossil contents were carried out on the ditch cuttings samples. These results were integrated to generate the various interpretations observed in the study area.

2.1 Nannoplankton Preparation and Analysis

The sample cuttings were prepared for calcareous nannofossils contents using standard preparatory techniques. A small portion of the sediments about 5g of the sample was gently crushed in a mortar and dispersed in distilled water in a glass. It was dried at 50°C and mounted on a glass slide with the Norland adhesive mounting medium. Species were identified using relevant catalogues and materials. Quantitative measures were employed to get the total number of individual taxa, percentage of relative abundance, diversity from each sample. Taxa recovered from this study were grouped to enable palaeocological deductions. paleocological affinities of known published palaeocological preferences of some selected nannofossil species were considered and correlated with present work.

3. Results and Discussion

3.1 Lithostratigraphic description

Based on the gamma ray log and petrographic study of the cuttings within the study intervals (10510-13790ft), the lithology constitutes sandstones, shale/Mudstone, sandy mudstone, shaly sand and argillaceous sandstone units. The shale units occur in significant thicker section at the upper part of the well (Fig. 3), which suggests a more marine and lower part a paralic depositional setting. The sands ranged from fine to medium grains, sub-angular to rounded while the shales were mostly grey to dark grey in colour.




Figure 3. Lithologic section of well AS-2, Niger delta.

3.2 Abundance, diversity and Species richness of Calcareous nannoplankton

The abundance presentation of semi quantitative abundance values showed thus: 0 - (absent); 1 species - (present); 1-2 species- (rare); 2-5 species- (common); 5-15 species (abundant) and super abundant (> 15 species), this explains the distribution pattern of the recoverable taxa (Figure 4, and the selected species illustrated in Fig. 5). Nannoplankton assemblages recovered show very poor to scanty preservation at the lower part of the study interval (see distribution chart Fig. 4), while towards the upper second half of the interval preservation was relatively common to abundant. Their abundance increased and preservation became better in samples of the early Oligocene than their counterparts from the late Eocene.




Figure 4: Nannofossil distribution of AS-2 well, Niger Delta.

From the sampled interval, 140 species belonging to 10 genera, were identified. The nannoplankton assemblage were dominated by Reticulofenestra both in abundance and diversity followed by Cyclicargolithus, Coccolithus, Sphenolithus, Helicosphaera and others as seen in Fig. 4 & 6.

Based on the percentage occurrence and composition of individual calcareous nannoplankton in the well, Reticulofenestra group occurred most in more than half of the samples studied (Fig. 5). However, identification of some taxa could not go beyond the genus level for some forms: Reticulofenestra sp. Interminate coccolithus, Pontosphaera sp, Helicosphaera sp and Micrantolithus sp these phenomena could be as a result of poor preservation due to dissolution, hence may not contribute any diagnostic information in the intervals.

Those depths with maximum abundance and diversity of nannofossils reflect the zones of condensed sections representing 32Ma and? 35Ma assemblages based on the peak and bioevent that existed within the interval of 10510ft and 12010ft respectively, describing low rates of sedimentation within the period.




Figure 5: Some selected calcareous nannofossils in the studied section

3.3 Biozonation and age

Two major biozones, NP 21 and NP 20 of late Eocene and early Oligocene were identified using the globally recognized calcareous nannofossil zonation scheme of [20] and the Niger Delta Nannozones, (Figure 4). The last occurrence of *Reticulofenestra umbilica* and *Chiasmolithus oamaruensis* are zonal markers for the two biozones according to [20], it was suggested to be the boundary between early Oligocene and late Eocene. Different calcareous nannofossil species appeared in these zones.

Zonal Description

Nannofossil zone: Zone NP 21

Stratigraphic interval: 10510 - 12010 ft

Age: Early Oligocene (Rupelian)

Comments: It is marked by the last occurrence of *Reticulofenestra umbilica* and last occurrence of *Chiasmolithus oamaruensis* which are zonal marker for nannofossil of early Oligocene [20]. The zone is also marked by the first occurrence of *Cycliargolithus abisectus*, occurrence of *Coccolithus pelagicus* which is an age marker and also age determined by the appearance of *Helicosphaera Compacta*, Down-Hole appearance of *Cyclicargolithus floridanus* and *Sphenolithus moriformis* and last appearance of *Cyclicargolithus abisectus*.

Base: The base of this zone is marked by last depth occurrence of *Reticulofenestra umbilica* and also the first appearance of *Reticulofenestra umbilica* and also *Micrantolithus* spp.

Top: is marked by the last occurrence of *Helicosphaera* and also with the down-hole occurrence of *Cyclicargolithus floridanus*, *Reticulofenestra bisectus*, *Sphenolithus moriformis* and also the disappearance of *Chiasmolithus oamaruensis*.

Zonal Description

Nannofossil zone: ?NP 20 and Older

Stratigraphic interval: 12010 - 13790 ft

Age: Late Eocene (Priabonian)

Comments: The biozone was not determined due to the scarcity of calcareous nannofossil, this zone can be described with the obvious reduction of nannofossil abundance; it is the older zone in the studied well. It is marked by the first appearance of *Helicosphaera compacta* and *Micrantolithus* spp, is marked by the Down Hole occurrence of *Cyclicargolithus floridanus*. It was also suggested by the last occurrence of *Reticulofenestra Umbilica* and last occurrence of *Chiasmolithus oamaruensis* that appeared in the upper zone NP 21.

Base: The base of this zone is marked by the last appearance of *Reticulofenestra* spp. Nannofossil species are not abundant here.

3.4 Paleocological preferences

The Paleocological preferences of the taxa in the study interval were mainly interpreted from previous works of calcareous nannofossils with known paleocological preferences documented in various literatures. The distribution pattern of calcareous nannofossils assemblages show fluctuations and changes towards sea surface temperature, salinity, and nutrient availability. A total of 140 taxa were identified and recorded; however, only few species or species groups are chosen for detailed paleocological discussion. Species are grouped on the basis of taxonomic and paleocological affinities in order to enhance a trend. Calcareous nannofossils examination in the present study interval indicate dominance of *Reticulofenestra spp* with percentage abundance of 49% compared to the total taxa. Followed by *Cyclicargolithus spp*, (percentage abundance of 17%), *Sphenolithus spp*, (percentage abundance of 12%), *Coccolithus spp*, (percentage abundance of 8%), the taxa with an average percentage abundance of less than 5% were *Helicosphaera spp* 4%, *Pontosphaera spp* 3%, *Braarudosphaera spp* 2%, *Chiasmolithus oamaruensis* 1% (as seen in Fig. 6).




Figure 6: Pie chart of percentage occurrence of recovered taxa groups

3.5 Cool water Nannofossils Taxa

The following species *Reticulofenestra spp*, *Cyclicargolithus spp*, *Coccolithus spp* and *Chiasmolithus spp* have been characterized as cool water (Fig. 7), eutrophic nutrient rich species, and indicators of upwelling areas for this present study [5], [21-24].

Reticulofenestra spp

The *Reticulofenestra* group is the most predominant species (Fig. 4 & 6) in the study interval, consisting of *Reticulofenestra bisecta*, *Reticulofenestra scrippsae*, *Reticulofenestra umbilica*, *Reticulofenestra umbilica* and *Reticulofenestra spp*.

The average abundance of the group increased towards early Oligocene. According to [5], [21-22]; [24], this species is referred to cool water conditions. Their abundance suggest cool, more eutrophic and nutrient rich episode in the study which corresponds to the global event in the Oligocene.

Cyclicargolithus spp

This group consist of *Cyclicargolithus abiesctus* and *Cyclicargolithus floridanus*. They have been inferred as cool water species by [7], [24] and in this study, though ecological preference remains ambiguous by some authors [12]. They are the next in abundance in the study interval with 17% (Fig. 6) and because of their relationship in abundance with *Reticulofenestra* within the Oligocene (Fig. 7). The FO *Cyclicargolithus spp* appeared at depth 13360ft, disappeared and at the boundary of Eocene –

Oligocene reappeared again (Figure 4). The abundance of the group contributes to the cool water eutrophic environment.

Coccolithus spp

This group consist of *Coccolithus pelagicus* and *Coccolithus miopelagicus* making up 8% in percentage abundance of the total assemblage in the study (Fig. 6). Its first occurrence in this study appears in zone NP21 though very scanty occurrence. Presently, this species is considered as a typical cool water form [24], [26-27] and eutrophic upwelling zones [28], [29].

Chiasmolithus spp

Chiasmolithus group has only *Chiasmolithus oamaruensis* which appeared in the upper zone of NP2 (Figure 4). It is well established as a cold-water taxon in [4, 5],[30-32] suggested that this taxon was adapted to eutrophic environments.




Figure 7: Relative abundance of ecologically significant nannofossils of Total warm and cold water species against depth and blue dashed line signifies the Eocene-Oligocene boundary

3.6 Warm water Nannofossils taxa

Sphenolithus spp, *Helicosphaera spp*, *Pontosphaera spp* are characterize as warm water (Fig. 7), oligotrophic species [3] and [33-37].

Sphenolithus spp

Only *Sphenolithus moriformis* is present in this group from the study section. It has been characterized as oligotrophic warm water taxon by many authors [34, 35]. In this study the relative abundance is less compared to the *Reticulofenestra* which indicate cooler waters, hence the prevailing condition is suggestive of cooler- water and more eutrophic environment during this period.

Helicosphaera spp

This group consists of *Helicosphaera catrei* and *Helicosphaera compact* [34] and others which could not be identified beyond genus level (Fig. 4) due to poor preservation in the studied intervals. The presence of this group indicates hemiplegic - shelf environment, moderate nutrient and turbulence in the surface waters [38, 39].

Pontosphaera spp

Major examples of this group include *Pontosphaera multipora* and *Pontosphaera spp* (Fig. 4). This group indicate more common near shore and warm water conditions. These changes correlate with global sea level curves discussed by others [35].

4. Conclusions

This study examined palaeocological preferences of calcareous nannoplankton assemblages of AS-2 well in the Niger delta. The following observations were noted.

1. Quantitatively, the nannofossil assemblage recorded from Well AS-2, Niger Delta were scanty to moderately preserved, with 140 species belonging to 10 genera.
2. The assemblage belongs to the late Eocene – Oligocene or precisely dated as Priabonian – Rupelian, only two major nannofossil zones (NP21 and NP20) belonging to the early Oligocene to late Eocene age was identified using the standard zonation schemes of [20] and [40] for the studied interval of the well.
3. Presence of the following species *Reticulofenestra spp*, *Cyclicargolithus spp*, *Coccolithus spp* and *Chiasmolithus spp* have been characterized as cool water, eutrophic nutrient rich species, and indicators of upwelling zones for this present study.
4. Records of *Sphenolithus spp*, *Helicosphaera spp*, *Pontosphaera spp* are characterize as warm water, oligotrophic environments.
5. The prevailing environmental condition in accordance with the palaeocological preferences studied suggest cooler, more eutrophic condition for the studied section.

Funding: Not applicable.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: Not applicable

Conflicts of Interest: The authors declare no conflict of interest

References

1. Lohmann, H., (1902). Die Coccolithophoridae, eine Monographie der Coccolithen bildenden Flagellaten, zugleich ein Beitrag zur Kenntnis des Mittelmeerauftriebs. *Arch Protistenkd* 1:89–165.
2. Boboye, O. A. and Adeleye, A.M., (2009). High Resolution Biostratigraphy of Early Pliocene- Late Miocene Calcareous Nannoplankton and Foraminiferal, Deep Offshore, Niger Delta, Nigeria, *European Journal of Scientific Research* ISSN 1450-216X,34(3):308-325.
3. Raffi, I., Backman, J., Zachos, J.C., Sluijs, A., (2009). The response of calcareous nannofossils assemblages to the Paleocene Eocene Thermal Maximum at the Walvis Ridge in the South Atlantic. *Mar. Micropaleontol.* 70: 201–212.
4. Bralower, T.J., (2002). Evidence of surface water oligotrophy during the Paleocene-Eocene thermal maximum: Nannofossil assemblage data from Ocean Drilling Program Site 690, Maud Rise, Weddell Sea, *Paleoceanography*, 17(2): 1023.
5. Manooseh, P. K., Fatemeh, H., Ebrahim, G., Mousavi-Harami, R., (2014). Biostratigraphy and Paleocological Study of Pabdeh Formation in Interior Fars, Zagros Basin, Iran, *Open Journal of Geology* 4: 571- 581.
6. Umoh, E.E., Ehinola, O. A. and Osterlof P. L., (2011). An assessment of abundance and distribution patterns of major Calcareous nannofossil as observed from Oligocene –Pliocene subsurface sediments of the Niger Delta. *Nigerian Association of Petroleum Explorationists Bulletin* 23(2): 39 –48.
7. Auer, G., Piller, W.E., Harzhauser, M., (2014). High-resolution calcareous nannoplankton palaeoecology as a proxy for small-scale environmental changes in the Early Miocene. *Mar Micropaleontol* 111: 53–65
8. Oretade, B. S., and Ali, C. A., (2021). Calcareous nannofloras in Western Lobe Offshore, Niger Delta: Eutrophication and climate change implications. *Malaysian Journal of Society and Space* 17: 274-287.

9. Erba, E., and Tremolada, F., (2004). Nannofossil carbonate fluxes during the Early Cretaceous: Phytoplankton response to nitrification episodes, atmospheric CO₂, and an oxia. *Paleoceanography* 19: 1–18.
10. Umoh, E.E., Oyafunke, O.A. and Ighodaro, E.J., 2019. Calcareous Nannoplankton analysis of well BL-2, Deepwater Offshore, Niger Delta Basin. *Journal of Mining and Geology* 55(2): 119–126.
11. Haq, U.Z.B., (1971). Paleogene Calcareous Nannoflora. Part IV: Paleogene nannoplankton biostratigraphy and evolutionary rates in Cenozoic calcareous nannoplankton: Stockholm *Contr. Geology* 25(4): 129-158.
12. Adegoke, O. S., Ajibola, S.O., Edet, J.J., Osterloff, P.L., Ogbonnaya, K.U., (2016). Cenozoic Foraminifera and Calcareous Nannofossil Biostratigraphy of the Niger Delta. 570p
13. Ola, P.S., (2018). Morphological Description of Calcareous Nannofossils Assemblage of a Middle-Miocene to Late-Miocene Section in the Niger-Delta, Nigeria. *Open Journal of Geology*, 8: 925-937. <https://doi.org/10.4236/ojg.2018.89055>
14. Umoh, E.E., (2020a): Nanno-Bioevents and Condensed Section of Well-BXW Offshore, Niger Delta, Nigeria. *International Journal of Scientific Engineering and Science*, 4(5): 1-7.
15. Umoh, E. E., (2020b): An Integrated Study: Biofacies analyses of sediments of Well B-1, Offshore, Niger Delta. *Iconic Research and Engineering Journals* 3(10):316–325.
16. Lucas F.A., Omodolor H.E. (2018). Palynofacies Analysis, Organic thermal maturation and source rock Evaluation of sedimentary succession from Oligocene to Early Miocene Age in X2 well, Greater Ughelli Depobelt, Niger Delta Basin, Nigeria. *Journal of Geosciences and Geomatics* 6(2):85- 93.
17. Doust, H and Omatsola, E., (1990). Niger Delta. In: Edwards, J. D. and Santogrossi, P. A. (Eds.), *Divergent/passive Margin Basins. American Association of Petroleum Geologists Bulletin* 48: 201-238
18. Short, A.J., and Stauble, K.L., (1967). Outline of the Geology of the Niger Delta. *American Association of Petroleum Geologists Memoir* 51: 761-779.
19. Avbovbo, A.A., (1978). Tertiary lithostratigraphy of Niger Delta. *American Association of Petroleum Geologists Bulletin* 62(2): 295-300.
20. Martini, E., (1971). Standard Tertiary and Quaternary calcareous nannoplankton zonation. In: Farinacci (eds), *Proceedings II planktonic conference, Roma, 1970*, 2:739-785.
21. Newsam, C., Bown, P.R., Wade, B.S., Jones, H.L., (2017). Muted calcareous nannoplankton response at the Middle/Late Eocene Turnover event in the western North Atlantic Ocean. *Newsl. Stratigr.* 50:297–309.
22. Persico, D. & Villa, G., (2004): Eocene-Oligocene calcareous nannofossils from Maud Rise and Kerguelen Plateau (Antarctica): Paleocological and paleoceanographic implications. *Mar. Micropaleontology* 52(4): 153 – 179.
23. Saeedeh, S., and Azam M., (2022). Calcareous nannofossils of the late Eocene- early Oligocene from the Pabdeh - Asmari transition in Dezful embayment (SW Iran): Evidence of a climate cooling event. *Paleogeography and Paleoclimatic journal* 2: 2-22.
24. Villa, G., Fioroni, C., Pea, L., Bohaty, S., Persico, D., (2008). Middle Eocene–late Oligocene climate variability: Calcareous nannofossils response at Kerguelen Plateau, Site 748. *Marine Micropaleontology* 69: 173–192
25. Fioroni, C., Villa, G., Persico, D., Jovane, L., (2015). Middle Eocene-Lower Oligocene calcareous nannofossil biostratigraphy and paleoceanographic implications from Site 711 (equatorial Indian Ocean). *Mar. Micropaleontol.* 118: 50–62.
26. Marino, M., Maiorano, P., Flower, B.P., (2011). Calcareous Nannofossil changes during the Mid-Pleistocene Revolution: Paleocologic and paleoceanographic evidence from North Atlantic Site 980/981. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 306:58–69.
27. Kameo, K., Kubota, Y., Haneda, Y., Suganuma, Y., and Okada, M., (2020). Calcareous nannofossil biostratigraphy of the Lower–Middle Pleistocene boundary of the GSSP, Chiba composite section in the Kokumoto Formation, Kazusa Group, central Japan, and implications for seasurface environmental changes, *Progress in Earth and Planetary Science*, 7: 36
28. Cachão, M., Moita, M.T., (2000). *Coccolithus pelagicus*, a productivity proxy related to moderate fronts off Western Iberia. *Mar. Micropaleontol.* 39: 131–155.
29. Tangunan, D.N., Baumann, K.-H., Just, J., LeVay, L.J., Barker, S., Brentegani, L., De Vleeschouwer, D., Hall, I.R., Hemming, S., Norris, R., (2018). The last 1 million years of the extinct genus *Discoaster*: Plio-Pleistocene environment and productivity at Site U1476 (Mozambique Channel). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 505: 187–197.
30. Wei, W. C., & Wise, S. W. (1990). Biogeographic Gradients of Middle Eocene Oligocene Calcareous Nannoplankton in the South-Atlantic Ocean. *Palaeogeography Palaeoclimatology Palaeoecology* 79(1–2):29–61.
31. Firth, J. V., and Wise, Jr., S. W., (1992). A preliminary study of the evolution of *Chiasmolithus* in the middle Eocene to Oligocene of Sites 647 and 748, ODP Leg 120. *Proc. Ocean Drill. Program Sci. Results* 120:493– 508.
32. Aubry, M.P., (1992). Paleogene Calcareous Nannofossils from the Kerguelen Plateau, Leg 120. In: Wise, S.W., Schlich, R., et al., Eds., *Proceedings of the Ocean Drilling Program, Scientific Results, Ocean Drilling Program, College Station*, 471-491.
33. Cao, W., Xi, D., Melinte-Dobrinescu, M.C., Jiang, T., Wise, S.W., Wan, X., (2018). Calcareous nannofossil changes linked to climate deterioration during the Paleocene-Eocene thermal maximum in Tarim Basin, NW China. *Geosci. Front.* 9: 1465–1478.
34. Gibbs, S.J., Bralower, T.J., Bown, P.R., Zachos, J.C., Bybell, L.M., (2006). Shelf and open ocean calcareous phytoplankton assemblages across the Paleocene-Eocene Thermal Maximum: implications for global productivity gradients. *Geology* 34: 233–236.
35. Švábenická, L., Bubik, M. and Stráňík, Z. (2007) Biostratigraphy and Paleoenvironmental Changes on the Transition from the Menilite to Krosno Lithofacies (Western Carpathians, Czech Republic). *Geology Carpathians* 58: 327-262.
36. Imai, R., Farida, M., Sato, T., Iryu, Y., (2015). Evidence for eutrophication in the northwestern Pacific and eastern Indian oceans during the Miocene to Pleistocene based on the nannofossil accumulation rate, *Discoaster* abundance, and coccolith size distribution of *Reticulofenestra*. *Mar. Micropaleontol.* 116: 15–27.
37. Kalb, A.L., Bralower, T.J., (2012). Nannoplankton origination events and environmental Changes in the late Paleocene and early Eocene. *Mar. Micropaleontol.* 93: 1–15.
38. Perch-Nielsen K., (1985b). Cenozoic calcareous nannofossils. In: Bolli HM, Saunders JB, Perch-Nielsen K (eds) *Plankton Stratigraphy*. Cambridge University Press, 427-554p

39. Flores, J. A., & Sierro, F. J., & Raffi, I., (1995). Evolution of the calcareous nannofossils assemblage as a response to the paleoceanographic changes in the eastern equatorial Pacific Ocean from 4 to 2 Ma. Leg 138, sites 849 and 852. In Pisias, N. G., Mayer, L. A., Janecek, T. R., Palmer-Julson, A., & van Andel, T. H. (Eds.), Proceedings of the ocean drilling program scientific results, College station TX 138: 163–176.

Cite article as:

Umoh, E.E. Palaeoecological Aspects of Nannoplankton Assemblages of AS-2 Well, Niger Delta. *Ajayi Crowther J. Pure Appl. Sci.* 2023, 2(2), pp. 107-117. | doi: <https://doi.org/10.56534/acjpas.2023.02.02.107>.